

Blood Pressure monitoring procedure for carer

Action	Rationale
Explain procedure; obtain and record verbal	To ensure that the resident is aware and
consent	consents to the procedure
Wash/use sanitizer on your hands	To avoid cross infection and contamination in
	line with infection control policies.
Position the residents/service user in a sitting position with their arm comfortably supported prior to taking the reading. In certain situations, you may be asked to do a sitting and standing blood pressure reading. If so, the cuff should be left in place while you assist the resident/service user to stand and then repeat the reading.	To obtain the most accurate result.
Ensure that the reading is done when the	To ensure that you have an accurate result as
resident/service user is relaxed and rested.	any stimulant or exertion can result in a higher
Avoid taking the reading soon after they have drunk coffee or other stimulant, or if after exertion.	reading than is normal for the resident/service user.
Turn blood pressure monitor on, look that it is	This check is to make sure that the batteries
working before you start the procedure.	and the blood pressure monitor is in good
working before you start the procedure.	working order. Replace batteries if necessary.
Apply a suitably sized cuff (e.g. if	To ensure that you get an accurate reading.
resident/service user is obese use a large cuff if	The choice and you got an accurate reading.
one is available) and check that it is	
comfortable. Explain that the cuff will become	
tighter for a short while to get a reading.	
Wash/use sanitizer on your hands.	To avoid cross infection and contamination
,	between staff/residents in line with infection
	control.
Document results in the service	To avoid error and communicate ensure safe
user's/resident's records/integrated care	accurate readings are entered in the correct
records immediately.	user record for the Care Home ready to relay to
	the requesting clinician.
Report the results to the appropriate Health	To ensure that the resident/service user
Care Professional as soon as possible in a pre-	receives prompt, appropriate care and if
agreed way e.g., email/phone/in person if they	necessary is on the correct dose of medication.
are visiting.	
Clean the blood pressure monitor and cuff with	T prevent any spread of infection and to ensure
anti-bacterial/sterile wipes.	that the device is ready and safe to use when next needed.
Store the blood pressure monitor/cuff in an	To save time when a request is made for a
easily accessible place where it can be found	blood pressure reading to be done. Ann
when needed.	